

1.2. High Performance Liquid Chromatography

1. Application of the Shats equation for prediction of retention values in reversed- and normal-phase liquid chromatography/ Grizodoub A.I., Levin M.G., Asmolova N.N., Georgievsky V.P. // Журн. аналит. химии. - 1992. - Т.47, № 3. - С.514-521.
2. Model of Mixed Adsorption Centre for Description of Retention in Liquid Chromatography with Multicomponent Mobile Phases/ Levin M.G., Grizodoub A.I., Asmolova N.N. et al.// Chromatographia. - 1993. -Vol.37, N 9/10. - P. 517-524.
3. About retention mechanism in reversed-phase liquid chromatography with multicomponent mobile phases /Levin M.G., Grizodoub A.I., Asmolova N.N. et al. // Журн. аналит. химии. - 1993. - Т.48, № 11. - С.81-93.
4. Levin M., Grizodoub A., Leontiev D., Gergievsky V. Retention in Liquid Chromatography with Multicomponent Mobile Phases: Relationship with Binary ones// Chromatographia. - 1995. - Vol. 40, N 5/6. P. - 321-328.
5. Kulikov A.U., Verushkin A.G., Shklyaeв S.A. High performance liquid chromatography in pharmaceutical analysis. Part 1. L1 columns or C18 silica bounded phases. About column interchangeability // Фармаком. – 2001. - № 1. – С. 47-56.
6. Kulikov A.U., Verushkin A.G. High performance liquid chromatography in pharmaceutical analysis. “System suitability” test // Вісник ХНУ. Серія «Хімія». – 2002. № 549. Вип. 8(31). – С. 57-65.
7. Kulikov A.U., Verushkin A.G. High performance liquid chromatography in pharmaceutical analysis. Part 2. About acceptable changes in chromatographic conditions // Фармаком. – 2002. - № 4. – С. 59-68.
8. Kulikov A.U. Column interchangeability in assays with use of reversed-phase high performance liquid chromatography // Фармаком. – 2008. - № 2. –С. 17-45.
9. Properties of 2,4-dinitrophenyl derivatives of amino acids as analytical forms for HPLC / Boichenko A.P., Chernyshova O.S., Kulikov A.Y., Loginova L.P. // Russ. J. Appl. Chem. – 2011. – Vol. 84, N 6. – P. 957-963.